

Accelerating Expansion of the Universe with Nonlinear Spinors

Ying-Qiu Gu*

Department of Mathematics, Fudan University, Shanghai, 200433, China

Abstract

The accelerating expansion of the present Universe is an exciting challenge to the standard cosmology, which reflects that our current theories on matter have some incompleteness. In this paper, we analyze the cosmological model with nonlinear spinor fields source in detail. The results may be able to provide a natural explanation for the puzzles of the acceleration and negative pressure.

PACS numbers: 98.80.Bp, 98.80.Cq, 98.80.Jk, 95.35.+d

Key Words: *accelerating expansion, negative pressure, nonlinear spinor, dark matter*

1 Introduction

The observational data show that the expansion of the present Universe is accelerating[1, 2, 3, 4]. The fundamental quest in theoretical cosmology is to identify the origin of this phenomenon. In principle there are two possibilities: (1) the presence of matter has negative pressure which violates the strong energy condition, (2) modified Friedmann equations[5]. Numerous researches along this direction come forth recently. The following work investigates the properties of a cosmological model with the dominant spinor fields as matter source, which is a further research of the preceding papers[6, 7, 8]. Some similar works were once provided in[9, 10, 11, 12], where the authors also considered the nonlinear spinor field as matter model, but only one spinor field was taken into account.

*email: yqgu@luody.com.cn

In this paper, we translate the dynamic variables of the spinor fields into the equivalent concepts of mass density ρ_m and pressure P of perfect fluid model, and express these variables as well as the other geometrical variables such as the expanding speed $\dot{a}$, the acceleration $\ddot{a}$, the Hubble's parameter H , as the functions of scale factor a , and then give detailed numerical computation. The influences of the parameters of this model on the behavior of the Universe are also considered. The results may be able to provide a natural explanation for the puzzles of acceleration and negative pressure.

2 Dynamic equation

For the Friedman-Robertson-Walker(FRW) metric

$$g_{\mu\nu} = a^2(t)\text{diag}[1, -1, -f^2(r), -f^2(r)\sin^2\theta], \quad (2.1)$$

with

$$f = \begin{cases} \sin r & \text{if } K = 1, \\ r & \text{if } K = 0, \\ \sinh r & \text{if } K = -1, \end{cases} \quad (2.2)$$

we derived the dynamic equation with nonlinear spinor source as follows[6]

$$a'^2 + Ka^2 - \frac{1}{3}\Lambda a^4 = \frac{8\pi G a}{3\Omega} \sum_{X_k \in \Omega} (\mathbf{K}_k + m_k) + C_0, \quad (2.3)$$

where $a' = \frac{d}{dt}a$, $d\Omega = f^2 \sin\theta dr d\theta d\varphi$ stands for the angular volume independent of t , C_0 is a negative integral constant closely relevant to P , and $\mathbf{K}_k$ is the kinetic energy of k -th spinor. In statistical sense we have the mean kinetic energy as follows[13]

$$\bar{\mathbf{K}} = \frac{3}{2}kT(a) = \frac{3\bar{m}b^2}{5a(a + \sqrt{a^2 + b^2})}, \quad (2.4)$$

with $\bar{m}$ being the mean mass of all particles, and b is a constant determined by the initial data

$$\frac{b}{a_0} = \sqrt{\frac{5kT_0}{\bar{m}} \left(1 + \frac{5kT_0}{4\bar{m}}\right)} \ll 1, \quad (1\text{eV} = 2.32 \times 10^4 K). \quad (2.5)$$

Detail analysis shows we have solution of physical meanings only if $K = 1$, i.e. the closed space.

Denote

$$R = \frac{4\pi G}{3\Omega} \sum_{X_k \in \Omega} m_k, \quad C_0 = -R^2(1 - \delta^2), \quad (0 < \delta < 1), \quad (2.6)$$

where R is a cosmic scale independent of a , δ is a dimensionless number relevant to pressure P . Substituting (2.4) and (2.6) into (2.3) we finally get

$$a'^2 = \frac{6}{5}R(\sqrt{a^2 + b^2} - a) + \delta^2 R^2 + \frac{1}{3}\Lambda a^4 - (a - R)^2, \quad (2.7)$$

we have singular-free cosmological model if

$$b < \frac{5}{6}R(1 - \delta^2). \quad (2.8)$$

For the extreme case with $b = \Lambda = 0$, we have the simple harmonic solution

$$a = R(1 - \delta \cos t), \quad f = \sin r. \quad (2.9)$$

Compared with observational data, a rough computation shows

$$\delta = 0.8 \sim 1, \quad R = (0.55 \sim 1.85) \times 10^{12} \text{yr}, \quad t_a = 20^\circ \sim 30^\circ \quad (2.10)$$

where t_a is the present time.

3 Functions and Properties

Theoretically, by (2.7) together with the concrete values of constants δ, b, λ , we can exactly determine the solution. But we will encounter with difficulties in practical calculation, because the solution of (2.7) can not expressed by elementary functions and the definite values of δ, b, λ are unknown. However the problem can be much simplified if we take a but not t as independent variable. In what follows, we adopt this method and make numerical computation for some important variables.

We take $R = 1$ as the length unit. Let $\lambda = \frac{1}{3}\Lambda R^4$, we get the dimensionless form of Eq.(2.7) as follows

$$a'^2 = \frac{6}{5}(\sqrt{a^2 + b^2} - a) + \delta^2 + \lambda a^4 - (a - 1)^2, \quad (3.1)$$

$$a'' = \frac{3}{5} \frac{a - \sqrt{a^2 + b^2}}{\sqrt{a^2 + b^2}} + 1 - a + 2\lambda a^3, \quad (3.2)$$

Let τ be the cosmic time, i.e. $d\tau = a dt$, then we have the following dimensionless expanding speed $\dot{a}$ and acceleration $\ddot{a}$ expressed by a

$$\dot{a} = \frac{da}{d\tau} = \frac{da}{adt} = \pm \frac{1}{a} \left(\frac{6}{5}(\sqrt{a^2 + b^2} - a) + \delta^2 + \lambda a^4 - (a - 1)^2 \right)^{\frac{1}{2}}, \quad (3.3)$$

$$\ddot{a} = \frac{d^2a}{d\tau^2} = \frac{1}{a^3} \left(-\frac{3}{5} \frac{(a^2 + 2b^2)}{\sqrt{a^2 + b^2}} - \frac{2}{5}a + \lambda a^4 + 1 - \delta^2 \right). \quad (3.4)$$

By (3.4) we have

$$\ddot{a} \rightarrow \frac{1}{a^3} \left(1 - \delta^2 - \frac{6}{5}b \right) \rightarrow +\infty, \quad \text{as } a \rightarrow 0.$$

So the Universe indeed has a period of inflation, the following numerical results clearly show this period. By (2.8), we find the condition of singular-free is just the condition of inflation.

Now we firstly translate the dynamic variables of the nonlinear spinor model (2.7) or (3.1) into the equivalent ρ_m, P, w of the usual fluid model via the Einstein equation, with which all scientists are more familiar, and then look for the relations among them and the variable a as well as the parameters δ, b, λ . By calculation we have the scalar curvature $\mathcal{R}$ and Einstein's tensor $G_{\mu\nu}$ as

$$\mathcal{R} = 6\frac{a'' + Ka}{a^3}, \quad G_{00} = -3\left(\frac{a'^2}{a^2} + K\right), \quad G_{11} = 2\frac{a''}{a} - \frac{a'^2}{a^2} + K, \quad (3.5)$$

and $G_{22} = G_{11}f^2$, $G_{33} = G_{11}f^2 \sin^2 \theta$. Substituting (3.5) into Einstein equation with fluid model

$$T_\nu^\mu = -\frac{1}{8\pi G}(G_\nu^\mu + \Lambda\delta_\nu^\mu) = (\rho_m + P)u^\mu u_\nu - P\delta_\nu^\mu, \quad (3.6)$$

for static fluid, we get

$$\rho_m = -\frac{1}{8\pi G}(G_0^0 + \Lambda) = \frac{1}{8\pi Ga^4}[3(a'^2 + Ka^2) - \Lambda a^4], \quad (3.7)$$

$$P = \frac{1}{24\pi G}[-(G_0^0 + \mathcal{R}) + 3\Lambda] = \frac{1}{8\pi Ga^4}(a'^2 - 2aa'' - Ka^2 + \Lambda a^4). \quad (3.8)$$

Define the density unit $\rho_0 = \frac{1}{8\pi GR^2} = 1$, and substituting the dynamic equation of spinor model (3.1) and (3.2) into ρ_m, P , then we get the dimensionless functions of the nonlinear spinor fields corresponding to the fluid model

$$\rho_m = \frac{3}{a^4} \left(\delta^2 - 1 + \frac{2}{5}(3\sqrt{a^2 + b^2} + 2a) \right), \quad (3.9)$$

$$P = \frac{1}{a^4} \left(\delta^2 - 1 + \frac{6}{5} \frac{b^2}{\sqrt{a^2 + b^2}} \right), \quad (3.10)$$

$$w = \frac{P}{\rho_m} = \frac{5\sqrt{a^2 + b^2}(\delta^2 - 1) + 6b^3}{3[6(a^2 + b^2) + \sqrt{a^2 + b^2}(5(\delta^2 - 1) + 4a)]}. \quad (3.11)$$

By the above equation, we find ρ_m, P, w are all continuous functions of a and constants δ, b , but independent of λ as they should be. The Hubble's parameter is defined by

$$H = \frac{da}{ad\tau} = \frac{a'}{a^2} = \frac{1}{a^2} \left(\frac{6}{5}(\sqrt{a^2 + b^2} - a) + \delta^2 + \lambda a^4 - (a - 1)^2 \right)^{\frac{1}{2}}. \quad (3.12)$$

In addition, the angular time t and the cosmic time τ can be expressed by the integrals of a

$$t = \int_{a_{min}}^a \frac{1}{a'} da, \quad \tau = \int_{a_{min}}^a \frac{a}{a'} da, \quad (3.13)$$

where a_{min} stands for the minimum of a . τH is also an important dimensionless variables. According to the functions defined above, we can easily compute the values, and clearly describe the properties of the corresponding cosmological model.

The numerical results are shown in following figures. Fig.(1), (2) and (3) display the phase trajectories of $a' \sim a$ with the change of parameters b, λ and δ respectively.




Figure 1: with varying b




Figure 2: with varying λ




Figure 3: with varying δ




Figure 4: Evolution curve & τH

By these figures we find that, b mainly influences the behavior as $a \rightarrow a_{min}$, in contrast, λ influences the behavior as $a \rightarrow a_{max}$. The influences of δ on a are more symmetrical.

By Fig.(4) we find the curve of $a(t)$ is quite near the simple harmonic function. The cosmic time τ increases slowly as $a \rightarrow a_{min}$, but more fast as $a \rightarrow a_{max}$. Except for the two extreme cases of a , we find the dimensionless function τH almost acts like a line for the most cases. The typical value is $0.5 < \tau H < 0.7$.

Fig.(5) and Fig.(6) show the Inflating and accelerating expanding process of the Universe near $a \rightarrow a_{min}$. In contrast, the cosmic time τ evolves very slowly in this period. These two factors mean that, the Universe just spends a very short time to complete her rebirth.

By Fig.(5) we find H varying slowly except for the case near a_{min} . This is the reason why the Hubble's law is quite effective.

Fig.(7) shows the evolution of the equivalent mass density ρ_m and pressure P of the spinor fields. In contrast with the fluid model, for the spinor fields we have two




Figure 5: The acceleration $\ddot{a}(\tau)$ & the Hubble's parameter H

distinct differences: (1) the pressure is negative, (2) the density ρ_m is not a monotone decreasing function of a . This means that, a spinor can not be simply treated as a dead point particle while the spacetime is heavily curved, which strongly interplays with the spacetime for this case. However it becomes normal as the spacetime become flat, e.g. as $a > 0.2$ shown by Fig.(7).

The curve of the equation of state $w = \frac{P}{\rho_m}$ is shown in Fig.(6), which seems to be quite coincident with the observational data.

4 Conclusion

Now we extract some important information from the above analysis and figures:

(C1). The cosmological factor Λ have little influence on the young Universe, and by Fig.(2) we find that Λ is purely superfluous without any effective use but increasing theoretical troubles, so setting $\Lambda = 0$ is a wise choice.

(C2). The temperature parameter b increases the singular as $a \rightarrow 0$, but has little influence on a normal evolving Universe. Noticing the small value of b , we find (2.9) gives a good approximation of the solution.

(C3). The Universe of nonlinear spinors has an inflating and accelerating expanding period in her early age. The inflation means the Universe only spends a very short time to get her rebirth and recover her vitality.




Figure 6: The expanding speed $\dot{a}(\tau)$ and the equation of state $w = \frac{P}{\rho_m}$




Figure 7: The relation between ρ_m, P and the scale factor a

(C4). Different from the fluid model, the spinor fields provide negative pressure, and the equivalent density ρ_m of the spinor fields is not a monotone decreasing function of a . Spinor strongly interplays with the heavily curved spacetime.

References

- [1] A. G. Riess *et al.* (Supernova Search Team), *Astron. J.* 116, 1009 (1998), arXiv:astro-ph/9805201.
- [2] S. Perlmutter *et al.* (Supernova Cosmology Project), *Astrophys. J.* 517, 565 (1999), arXiv:astro-ph/9812133.
- [3] N. Spergel *et al.* (WMAP), *Astrophys. J. Suppl.* 148, 175 (2003), arXiv:astro-ph/0302209.
- [4] M. Tegmark *et al.* (SDSS), *Phys. Rev. D* 69, 103501 (2004), arXiv:astro-ph/0310723.
- [5] Marek Szydlowski, *Cosmological zoo C accelerating models with dark energy*, arXiv:astro-ph/0610250
- [6] Y. Q. Gu, *A Cosmological Model with Dark Spinor Source*, arXiv:gr-qc/0610147.
- [7] Y. Q. Gu, *Some properties of the spinor soliton*, *Adv. appl. Cliff. Alg.* **8(1)**, 17-29(1998), <http://www.clifford-algebras.org/v8/81/gu81.pdf>
- [8] Y. Q. Gu, *Characteristic Functions and Typical Values of the Nonlinear Dark Spinor*, arXiv:hep-th/0611210
- [9] Saha, Bijan, *Phys. Rev. D* **64** (2001) 123501.
- [10] Saha, Bijan and Boyadjiev, T., *Phys. Rev. D* **69** (2004) 124010.
- [11] Bijan Saha, *Nonlinear spinor field in Bianchi type-I cosmology: accelerated regimes*, arXiv:gr-qc/0608047
- [12] Ribas, M.O., Devecchi, F.P., and Kremer, G.M., *Phys. Rev. D* **72**, (2005)123502, arXiv:gr-qc/0511099.
- [13] Y. Q. Gu, G. J. Ni, *The Relation between Cosmic Temperature and Scale Factor*, arXiv:gr-qc/0609098.